

FORM PTO-1390
(REV 10-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C.371****GIN-6713CPUS**

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/554933

INTERNATIONAL APPLICATION

PCT/JP98/05238

INTERNATIONAL FILING DATE

20 November 1998 (20.11.98)

PRIORITY DATE CLAIMED

25 November 1997 (25.11.97)

TITLE OF INVENTION

**HUMAN PROTEINS HAVING TRANSMEMBRANE DOMAINS AND DNAs ENCODING
THESE PROTEINS**

APPLICANT(S) FOR DO/EO/US

Seishi KATO; Tomoko KIMURA; and Shingo SEKINE

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C.371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371 (f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371 (b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)) (**unexecuted**) (**5 sheets**);
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included
13. ☐ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: **Transmittal Letter (2 sheets in duplicate); Sequence Diskette Transmittal Letter with Sequence Listing Diskette and hard copy of Sequence Listing attached (26 sheets); PCT Notification of Receipt of Record Copy (PCT/IB/301) (2 sheets); PCT Notification Concerning Submission or Transmittal of Priority Document (Form PCT/IB/304) (1 sheet); PCT Notification of the Recording of a Change (Form PCT/IB/306) (1 sheet); PCT International Published Application (WO 99/27094) (without International Search Report) (89 sheets); PCT International Search Report (Form PCT/ISA/210) (5 sheets); PCT Information Concerning Elected Offices Notified of their Election (Form PCT/IB/332) (1 sheet); Cover Page of PCT International Published Application (WO 99/27094) (1 sheet); PCT Notification of Transmittal of the International Preliminary Examination Report (Form PCT/IPEA/416) (9 sheets); PCT Written Opinion (7 sheets); Check (#037073) (\$1100); Certificate of Express Mailing (1 sheet); and Return Postcard.**

COPY

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

09/554933

INTERNATIONAL APPLICATION NO.

PCT/JP98/05238

ATTORNEY'S DOCKET NO.

GIN-6713CPUS

17. ☒ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) .(a/o January 1, 2000):**

Search Report has been prepared by the EPO or JPO.....\$840
 International preliminary examination fee paid to
 USPTO (37 CFR 1.482).....\$670
 No international preliminary examination fee paid to
 USPTO (37 CFR 1.482) but international search fee
 paid to USPTO (37 CFR 1.445(a)(2)).....\$690
 Neither international preliminary examination fee
 (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2))
 paid to USPTO.....\$670
 International preliminary examination fee paid to
 USPTO (37 CFR 1.482) and all claims satisfied provisions
 of PCT Article 33(2)-(4).....\$96

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$840

Surcharge of \$130.00 for furnishing the oath or declaration later than 20 30
 months from the earliest claimed priority date (37 CFR 1.492(e)).

\$--

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	11 -20 =	0	X \$18.00
Independent claims	2 -3 =	0	X \$78.00

\$0

\$0

MULTIPLE DEPENDENT CLAIM(S) (if applicable)

+ 260.00

\$260

TOTAL OF ABOVE CALCULATIONS =

\$1100

Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity
 Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28)

\$--

SUBTOTAL =

\$1100

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
 months from the earliest claimed priority date (37 CFR 1.492(f)).

\$--

TOTAL NATIONAL FEE =

\$1100

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment
 must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31).
 \$40.00 per property

\$--

TOTAL FEES ENCLOSED =

\$1100

Amount to be:
refunded

\$

charged

\$

a. ☒ A check in the amount of \$1100 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees.
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit
 any overpayment to Deposit Account No. 12-0080. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Amy E. Mandragouras, Esq.
LAHIVE & COCKFIELD, LLP
28 State Street
Boston, Massachusetts 02109
United States of America
(617)227-7400
Date: 22 May 2000

SIGNATURE

Peter C. Lauro

NAME

32,360

REGISTRATION NUMBER

3/PR+5

1

526 Rec'd FCT/PTO 22 MAY 2000

DESCRIPTION

Human Proteins Having Transmembrane Domains and DNAs Encoding these Proteins

5

TECHNICAL FIELD

The present invention relates to human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells expressing said cDNAs. The proteins of the present invention can be employed as pharmaceuticals or as antigens for preparing antibodies against said proteins. The human cDNAs of the present invention can be utilized as probes for the gene diagnosis and gene sources for the gene therapy. Furthermore, the cDNAs can be utilized as gene sources for large-scale production of the proteins encoded by said cDNAs. Cells, wherein these membrane protein genes are introduced and membrane proteins are expressed in large amounts, can be utilized for detection of the corresponding ligands, screening of novel low-molecular pharmaceuticals, and so on.

BACKGROUND ART

Membrane proteins play important roles, as signal
25 receptors, ion channels, transporters, etc. in the material
transportation and the information transmission which are
mediated by the cell membrane. Examples thereof include
receptors for a variety of cytokines, ion channels for the
sodium ion, the potassium ion, the chloride ion, etc.,
30 transporters for saccharides and amino acids, and so on,
where the genes of many of them have been cloned already.

It has been clarified that abnormalities of these membrane proteins are associated with a number of hitherto-cryptogenic diseases. For instance, a gene of a membrane protein having twelve transmembrane domains was identified as the gene responsible for cystic fibrosis [Rommens, J. M. et al., Science 245: 1059-1065 (1989)]. In addition, it has been clarified that several membrane proteins act as receptors when a virus infects the cells. For instance, HIV-1 is revealed to infect into the cells through mediation of a membrane protein fusin having a membrane protein on the T-cell membrane, a CD-4 antigen, and seven transmembrane domains [Feng, Y. et al., Science 272: 872-877 (1996)]. Therefore, discovery of a new membrane protein is anticipated to lead to elucidation of the causes of many diseases, so that isolation of a new gene coding for the membrane protein has been desired.

Heretofore, owing to difficulty in the purification, many membrane proteins have been isolated by an approach from the gene side. A general method is the so-called expression cloning which comprises transfection of a cDNA library in eucaryotic cells to express cDNAs and then detection of the cells expressing the target membrane protein on the membrane by an immunological technique using an antibody or a physiological technique on the change in the membrane permeability. However, this method is applicable only to cloning of a gene of a membrane protein with a known function.

In general, membrane proteins possess hydrophobic transmembrane domains inside the proteins, wherein, after synthesis thereof in the ribosome, these domains remain in

the phospholipid membrane to be trapped in the membrane. Accordingly, the evidence of the cDNA for encoding the membrane protein is provided by determination of the whole base sequence of a full-length cDNA followed by detection
5 of highly hydrophobic transmembrane domains in the amino acid sequence of the protein encoded by said cDNA.

DISCLOSURE OF INVENTION

The object of the present invention is to provide
10 novel human proteins having transmembrane domains, DNAs coding for said proteins, and expression vectors of said cDNAs as well as transformation eucaryotic cells that are capable of expressing said cDNAs.

As the result of intensive studies, the present
15 inventors have been successful in cloning of cDNAs coding for proteins having transmembrane domains from the human full-length cDNA bank, thereby completing the present invention. In other words, the present invention provides human proteins having transmembrane domains, namely
20 proteins containing any of the amino acid sequences represented by Sequence Nos. 1 to 3. Moreover, the present invention provides DNAs coding for the above-mentioned proteins, exemplified by cDNAs containing any of the base sequences represented by Sequence Nos. 4 to 6, 7, 9 and 11,
25 as well as transformation eucaryotic cells that are capable of expressing said cDNAs.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1: A figure depicting the
30 hydrophobicity/hydrophilicity profile of the protein encoded by clone HP01207.

Figure 2: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP01862.

Figure 3: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10493.

BEST MODE FOR CARRYING OUT THE INVENTION

The proteins of the present invention can be obtained, for example, by a method for isolation from human organs, cell lines, etc., a method for preparation of peptides by the chemical synthesis, or a method for production with the recombinant DNA technology using the DNAs coding for the transmembrane domains of the present invention, wherein the method for obtainment by the recombinant DNA technology is employed preferably. For instance, in vitro expression of the proteins can be achieved by preparation of an RNA by in vitro transcription from a vector having one of cDNAs of the present invention, followed by in vitro translation using this RNA as a template. Also, recombination of the translation region into a suitable expression vector by the method known in the art leads to production of a large amount of the encoded protein by using prokaryotic cells such as *Escherichia coli*, *Bacillus subtilis*, etc., and eucaryotic cells such as yeasts, insect cells, mammalian cells, etc.

In the case in which a protein of the present invention is produced by expression of one of the DNAs by in vitro expression, recombination of the translation region in said cDNA into a vector having an RNA polymerase

promoter, followed by addition into an in vitro translation system such as a rabbit reticulocyte lysate, a wheat germ extract, or the like, allows in vitro production of the protein of the present invention. Examples of the RNA polymerase promoter include T7, T3, SP6, and so on. Vectors containing such an RNA polymerase promoter are exemplified by pKA1, pCDM8, pT3/T7 18, pT7/3 19, pBluescript II, and so on. Also, addition of the dog pancreas microsome etc. in the reaction system enables the membrane protein of the present invention to be expressed in a form integrated in the microsome membrane.

In the case in which a protein of the present invention is produced by expression of a DNA in a microorganism such as *Escherichia coli* etc., a recombinant expression vector bearing the translation region in the cDNA of the present invention is constructed in an expression vector having an origin, a promoter, a ribosome-binding site, a cDNA-cloning site, a terminator etc., which can be replicated in the microorganism, and, after transformation of the host cells with said expression vector, the thus-obtained transformant is incubated, whereby the protein encoded by said cDNA can be produced on a large scale in the microorganism. In this case, a protein fragment containing an optional region can be obtained by carrying out the expression with inserting an initiation codon and a termination codon in front of and behind an optional translation region. Alternatively, a fusion protein with another protein can be expressed. Only a protein portion coding for said cDNA can be obtained by cleavage of said fusion protein with a suitable protease.

Examples of the expression vector for *Escherichia coli* include the pUC system, pBluescript II, the pET expression system, the pGEX expression system, and so on.

5 In the case in which one of the proteins of the present invention is produced by expression of a DNA in eucaryotic cells, the protein of the present invention can be produced as a transmembrane protein on the cell-membrane surface, when the translation region of said cDNA is subjected to recombination to an expression vector for
10 eucaryotic cells that has a promoter, a splicing region, a poly(A) insertion site, etc., followed by introduction into the eucaryotic cells. The expression vector is exemplified by pKA1, pED6dpc2, pCDM8, pSVK3, pMSG, pSVL, pBK-CMV, pBK-RSV, EBV vector, pRS, pYES2, and so on. Examples of
15 eucaryotic cells to be used in general include mammalian culture cells such as simian kidney cells COS7, Chinese hamster ovary cells CHO, etc., budding yeasts, fission yeasts, silkworm cells, *Xenopus laevis* egg cells, and so on, but any eucaryotic cells may be used, provided that they
20 are capable of expressing the present proteins on the membrane surface. The expression vector can be introduced in the eucaryotic cells by methods known in the art such as the electroporation method, the potassium phosphate method, the liposome method, the DEAE-dextran method, and so on.

25 After one of the proteins of the present invention is expressed in prokaryotic cells or eucaryotic cells, the objective protein can be isolated from the culture and purified by a combination of separation procedures known in the art. Such examples include treatment with a denaturing
30 agent such as urea or a surface-active agent, sonication,

enzymatic digestion, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity chromatography, reverse phase chromatography, and so on.

The proteins of the present invention include peptide fragments (more than 5 amino acid residues) containing any partial amino acid sequence in the amino acid sequences represented by Sequence Nos. 1 to 3. These peptide fragments can be utilized as antigens for preparation of antibodies. Hereupon, among the proteins of the present invention, those having the signal sequence are secreted in the form of maturation proteins on the surface of the cells, after the signal sequences are removed. Therefore, these maturation proteins shall come within the scope of the present invention. The N-terminal amino acid sequences of the maturation proteins can be easily identified by using the method for the cleavage-site determination in a signal sequence [Japanese Patent Kokai Publication No. 1996-187100]. Furthermore, some membrane proteins undergo the processing on the cell surface to be converted to the secretory forms. Such proteins or peptides in the secretory forms shall come within the scope of the present invention. When sugar chain-binding sites are present in the amino acid sequences, expression in appropriate eucaryotic cells affords proteins wherein sugar chains are added. Accordingly, such proteins or peptides wherein sugar chains are added shall come within the scope of the present invention.

The DNAs of the present invention include all DNAs

coding for the above-mentioned proteins. Said DNAs can be obtained by using a method by chemical synthesis, a method by cDNA cloning, and so on.

The cDNAs of the present invention can be cloned, for example, from cDNA libraries of the human cell origin. These cDNA are synthesized by using as templates poly(A)⁺ RNAs extracted from human cells. The human cells may be cells delivered from the human body, for example, by the operation or may be the culture cells. The cDNAs can be synthesized by using any method selected from the Okayama-Berg method [Okayama, H. and Berg, P., Mol. Cell. Biol. 2: 161-170 (1982)], the Gubler-Hoffman method [Gubler, U. and Hoffman, J. Gene 25: 263-269 (1983)], and so on, but it is preferred to use the capping method [Kato, S. et al., Gene 150: 243-250 (1994)], as exemplified in Examples, in order to obtain a full-length clone in an effective manner. In addition, commercially available, human cDNA libraries can be utilized. Cloning of the cDNAs of the present invention from the cDNA libraries can be carried out by synthesis of an oligonucleotide on the basis of an optional portion in the cDNA base sequences of the present invention, followed by screening using this oligonucleotide as the probe according to the colony or plaque hybridization by a method known in the art. In addition, the cDNA fragments of the present invention can be prepared by synthesis of an oligonucleotide to be hybridized at both termini of the objective cDNA fragment, followed by the usage of this oligonucleotide as the primer for the RT-PCR method from an mRNA isolated from human cells.

The cDNAs of the present invention are characterized

by containing either of the base sequences represented by Sequence Nos. 4 to 6 or the base sequences represented by Sequence Nos. 7, 9 and 11. Table 1 summarizes the clone number (HP number), the cells affording the cDNA, the total
5 base number of the cDNA, and the number of the amino acid residues of the encoded protein, for each of the cDNAs.

Table 1

Sequence No.	HP No.	Cell	Number of bases	Number of amino acids
1, 4, 7	HP01207	Stomach Cancer	2938	269
15 2, 5, 8	HP01862	Stomach Cancer	2290	311
3, 6, 9	HP10493	PMA-U937	3705	383

Hereupon, the same clones as the cDNAs of the present
20 invention can be easily obtained by screening of the cDNA libraries constructed from the human cell lines and human tissues utilized in the present invention by the use of an oligonucleotide probe synthesized on the basis of the cDNA base sequence described in any of Sequence Nos. 4 to 6, 7,
25 9 and 11.

In general, the polymorphism due to the individual difference is frequently observed in human genes. Accordingly, any cDNA that is subjected to insertion or deletion of one or plural nucleotides and/or substitution
30 with other nucleotides in Sequence Nos. 4 to 6, 7, 9 and 11 shall come within the scope of the present invention.

In a similar manner, any protein that is formed by these modifications comprising insertion or deletion of one

or plural amino acids and/or substitution with other amino acids shall come within the scope of the present invention, as far as the protein possesses the activity of any protein having the amino acid sequences represented by Sequence Nos. 1 to 3.

The cDNAs of the present invention include cDNA fragments (more than 10 bp) containing any partial base sequence in the base sequences represented by Sequence Nos. 4 to 6 or in the base sequences represented by Sequences No. 7, 9 and 11. Also, DNA fragments consisting of a sense chain and an anti-sense chain shall come within this scope. These DNA fragments can be utilized as the probes for the gene diagnosis.

In addition to the activities and uses described above, the polynucleotides and proteins of the present invention may exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

Research Uses and Utilities

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a

particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in

biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In

such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing

Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon γ , Schreiber, R.D. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6-Nordan, R. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and

Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

Immune Stimulating or Suppressing Activity

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders

(including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These
5 immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the
10 present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the
15 immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic
20 lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present
25 invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation),
30 may also be treatable using a protein of the present

invention.

Using the proteins of the invention it may also be possible to immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune

reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins in vivo as described in Lenschow et al., Science 257:789-792 (1992) and Turka et

al., Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function in vivo on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells in vitro with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the in vitro activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells in vivo.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity.

Tumor cells (e.g., sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected ex vivo with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection in vivo.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I α chain protein and β_2 microglobulin protein or an MHC class II α chain protein and an MHC class II β chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a

peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J.

Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

5 Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J.J. and Brunswick, M. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 10 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro 15 assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, 25 among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 30 1995;

182:255-260, 1995; Nair et al., Journal of Virology
67:4062-4069, 1993; Huang et al., Science 264:961-965,
1994; Macatonia et al., Journal of Experimental Medicine
169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical
5 Investigation 94:797-807, 1994; and Inaba et al., Journal
of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will
identify, among others, proteins that prevent apoptosis
after superantigen induction and proteins that regulate
10 lymphocyte homeostasis) include, without limitation, those
described in: Darzynkiewicz et al., Cytometry 13:795-808,
1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca
et al., Cancer Research 53:1945-1951, 1993; Itoh et al.,
Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology
15 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897,
1993; Gorczyca et al., International Journal of Oncology
1:639-648, 1992.

Assays for proteins that influence early steps of T-
cell commitment and development include, without limitation,
20 those described in: Antica et al., Blood 84:111-117, 1994;
Fine et al., Cellular Immunology 155:111-122, 1994; Galy et
al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad
Sci. USA 88:7548-7551, 1991.

Hematopoiesis Regulating Activity

25 A protein of the present invention may be useful in
regulation of hematopoiesis and, consequently, in the
treatment of myeloid or lymphoid cell deficiencies. Even
marginal biological activity in support of colony forming
cells or of factor-dependent cell lines indicates
30 involvement in regulating hematopoiesis, e.g. in supporting

the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelosuppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either in-vivo or ex-vivo (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In Culture of Hematopoietic Cells. R.I.

Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

Tissue Growth Activity

5 A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

10 A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well
15 as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic
20 plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-
25 forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue
30 destruction (collagenase activity, osteoclast activity,

etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors ex vivo for return in vivo to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as

a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium),

muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

Activin/Inhibin Activity

A protein of the present invention may also exhibit

activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins and are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin α family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- β group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population

to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W.Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

Hemostatic and Thrombolytic Activity

A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-

474, 1988.

Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-

1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160
1989; Stoltenborg et al., J. Immunol. Methods 175:59-68,
1994; Stitt et al., Cell 80:661-670, 1995.

Anti-Inflammatory Activity

5 Proteins of the present invention may also exhibit
anti-inflammatory activity. The anti-inflammatory activity
may be achieved by providing a stimulus to cells involved
in the inflammatory response, by inhibiting or promoting
10 cell-cell interactions (such as, for example, cell
adhesion), by inhibiting or promoting chemotaxis of cells
involved in the inflammatory process, inhibiting or
promoting cell extravasation, or by stimulating or
suppressing production of other factors which more directly
15 inhibit or promote an inflammatory response. Proteins
exhibiting such activities can be used to treat
inflammatory conditions including chronic or acute
conditions), including without limitation inflammation
associated with infection (such as septic shock, sepsis or
20 systemic inflammatory response syndrome (SIRS)), ischemia-
reperfusion injury, endotoxin lethality, arthritis,
complement-mediated hyperacute rejection, nephritis,
cytokine or chemokine-induced lung injury, inflammatory
bowel disease, Crohn's disease or resulting from over
25 production of cytokines such as TNF or IL-1. Proteins of
the invention may also be useful to treat anaphylaxis and
hypersensitivity to an antigenic substance or material.

Tumor Inhibition Activity

In addition to the activities described above for
immunological treatment or prevention of tumors, a protein

of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via ADCC). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth

Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive

disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

Examples

The present invention is embodied in more detail by the following examples, but this embodiment is not intended to restrict the present invention. The basic operations and the enzyme reactions with regard to the DNA recombination are carried out according to the literature ["Molecular Cloning. A Laboratory Manual " , Cold Spring Harbor Laboratory, 1989]. Unless otherwise stated, restrictive enzymes and a variety of modification enzymes to be used were those available from TAKARA SHUZO. The manufacturer's instructions were used for the buffer compositions as well as for the reaction conditions, in each of the enzyme reactions. The cDNA synthesis was carried out according to the literature [Kato, S. et al., Gene 150: 243-250 (1994)].

(1) Preparation of Poly(A)⁺ RNA

The histiocyte lymphoma cell line U937 (ATCC CRL 1593) stimulated by phorbol ester and tissues of stomach cancer delivered by the operation were used for human cells to extract mRNAs. The cell line was incubated by a conventional procedure.

After about 1 g of the human cells was homogenized in 20 ml of a 5.5 M guanidinium thiocyanate solution, a total mRNA was prepared according to the literature [Okayama, H. et al., "Method in Enzymology", Vol. 164, Academic Press, 1987]. This was subjected to chromatography on oligo(dT)-cellulose column washed with a 20 mM Tris-hydrochloride buffer solution (pH 7.6), 0.5 M NaCl, and 1 mM EDTA to obtain a poly(A)⁺ RNA according to the above-described literature.

(2) Construction of cDNA Library

Ten micrograms of the above-mentioned poly(A)⁺ RNA were dissolved in a 100 mM Tris-hydrochloride buffer solution (pH 8), one unit of an RNase-free, bacterial alkaline phosphatase was added, and the reaction was run at 37 °C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 50 mM sodium acetate (pH 6), 1 mM EDTA, 0.1% 2-mercaptoethanol, and 0.01% Triton X-100. Thereto was added one unit of a tobacco-origin acid pyrophosphatase (Epicentre Technologies) and a total 100 µl volume of the resulting mixture was reacted at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in water to obtain a solution of a decapped

poly(A)⁺ RNA.

The decapped poly(A)⁺ RNA and 3 nmol of a chimeric DNA-RNA oligonucleotide (5'-dG-dG-dG-dG-dA-dA-dT-dT-dC-dG-dA-G-G-A-3') were dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 7.5), 0.5 mM ATP, 5 mM MgCl₂, 10 mM 2-mercaptoethanol, and 25% polyethylene glycol, where to was added 50 units of T4RNA ligase and a total 30 µl volume of the resulting mixture was reacted at 20 °C for 12 hours. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in water to obtain a chimeric-oligo-capped poly(A)⁺ RNA.

After digestion of vector pKA1 (Japanese Patent Kokai Publication No. 1992-117292) developed by the present inventors with KpnI, about 60 dT tails were added using a terminal transferase. A vector primer to be used below was prepared by digestion of this product with EcoRV to remove a dT tail at one side.

After 6 µg of the previously-prepared chimeric-oligo-capped poly(A)⁺ RNA was annealed with 1.2 µg of the vector primer, the resulting product was dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 8.3), 75 mM KCl, 3 mM MgCl₂, 10 mM dithiothreitol, and 1.25 mM dNTP (dATP + dCTP + dGTP + dTTP), 200 units of a reverse transcriptase (GIBCO-BRL) were added, and the reaction in a total 20 µl volume was run at 42°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 7.5), 100 mM NaCl, 10 mM MgCl₂, and 1

mm dithiothreitol. Thereto were added 100 units of EcoRI and a total 20 μ l volume of the resulting mixture was reacted at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 20 mM Tris-hydrochloride buffer solution (pH 7.5), 100 mM KCl, 4 mM MgCl₂, 10 mM (NH₄)₂SO₄, and 50 μ g/ml of the bovine serum albumin. Thereto were added 60 units of an *Escherichia coli* DNA ligase and the resulting mixture was reacted at 16°C for 16 hours. To the reaction solution were added 2 μ l of 2 mM dNTP, 4 units of *Escherichia coli* DNA polymerase I, and 0.1 unit of *Escherichia coli* RNase H and the resulting mixture was reacted at 12°C for one hour and then at 22°C for one hour.

Next, the cDNA-synthesis reaction solution was used for transformation of *Escherichia coli* DH12S (GIBCO-BRL). The transformation was carried out by the electroporation method. A portion of the transformant was sprayed on the 2xYT agar culture medium containing 100 μ g/ml ampicillin and the mixture was incubated at 37°C overnight. A colony formed on the agar medium was picked up at random and inoculated on 2 ml of the 2xYT culture medium containing 100 μ g/ml ampicillin. After incubation at 37°C overnight, the culture mixture was centrifuged to separate the mycelia, from which a plasmid DNA was prepared by the alkaline lysis method. The plasmid DNA was subjected to double digestion with EcoRI and NotI, followed by 0.8% agarose gel electrophoresis, to determine the size of the cDNA insert. Furthermore, using the thus-obtained plasmid as a template, the sequence reaction was carried out by using an M13

universal primer labeled with a fluorescent dye and a Taq polymerase (a kit of Applied Biosystems) and then the product was examined with a fluorescent DNA sequencer (Applied Biosystems) to determine an about 400-bp base sequence at the 5'-terminus of the cDNA. The sequence data were filed as the homo/protein cDNA bank database.

(3) Selection of cDNAs Encoding Proteins Having Transmembrane Domains

A base sequence registered in the homo/protein cDNA bank was converted to three frames of amino acid sequences and the presence or absence of an open reading frame (ORF) beginning from the initiation codon was examined. Then, the selection was made for the presence of a signal sequence that is characteristic to a secretory protein at the N-terminus of the portion encoded by the ORF. These clones were sequenced from the both 5' and 3' directions by the use of the deletion method using exonuclease III to determine the whole base sequence. The hydrophobicity/hydrophilicity profiles were obtained for proteins encoded by the ORF by the Kyte-Doolittle method [Kyte, J. & Doolittle, R. F., J. Mol. Biol. 157: 105-132 (1982)] to examine the presence or absence of a hydrophobic region. In the case in which there is a hydrophobic region of a putative transmembrane domain in the amino acid sequence of an encoded protein, this protein was judged as a membrane protein.

(4) Functional Verification of Secretory Signal Sequence or Transmembrane Domains

It was verified by the method described in the literature [Yokoyama-Kobayashi, M. et al., Gene 163: 193-

196 (1995)] that the N-terminal hydrophobic region in the secretory protein clone candidate obtained in the above-mentioned steps functions as a secretory signal sequence. First, the plasmid containing the target cDNA was cleaved at an appropriate restriction enzyme site existing at the downstream of the portion expected for encoding the secretory signal sequence. In the case in which this restriction site was a protruding terminus, the site was blunt-ended by the Klenow treatment or treatment with the mung-bean nuclease. Digestion with HindIII was further carried out and a DNA fragment containing the SV40 promoter and a cDNA encoding the secretory signal sequence at the downstream of the promoter was separated by agarose gel electrophoresis. The resulting fragment was inserted between HindIII in pSSD3 (DDBJ/EMBL/GenBank Registration No. AB007632) and a restriction enzyme site selected so as to match with the urokinase-coding frame, thereby constructing a vector expressing a fusion protein of the secretory signal sequence of the target cDNA and the urokinase protease domain.

After *Escherichia coli* (host: JM109) bearing the fusion-protein expression vector was incubated at 37°C for 2 hours in 2 ml of the 2xYT culture medium containing 100 µg/ml of ampicillin, the helper phage M13K07 (50 µl) was added and the incubation was continued at 37°C overnight. A supernatant separated by centrifugation underwent precipitation with polyethylene glycol to obtain single-stranded phage particles. These particles were suspended in 100 µl of 1 mM Tris-0.1 mM EDTA, pH 8 (TE). Also, there were used as controls suspensions of single-stranded phage

particles prepared in the same manner from pSSD3 and from the vector pKA1-UPA containing a full-length cDNA of urokinase [Yokoyama-Kobayashi, M. et al., Gene 163: 193-196 (1995)].

5 The culture cells originating from the simian kidney, COS7, were incubated at 37°C in the presence of 5% CO₂ in the Dulbecco's modified Eagle's culture medium (DMEM) containing 10% fetal calf albumin. Into a 6-well plate (Nunc Inc., 3 cm in the well diameter) were inoculated 1 ×
10 10⁵ COS7 cells and incubation was carried out at 37°C for 22 hours in the presence of 5% CO₂. After the culture medium was removed, the cell surface was washed with a phosphate buffer solution and then washed again with DMEM containing 50 mM Tris-hydrochloric acid (pH 7.5) (TDMEM).
15 To the resulting cells was added a suspension of 1 µl of the single-stranded phage suspension, 0.6 ml of the DMEM culture medium, and 3 µl of TRANSFECTAM™ (IBF Inc.) and the resulting mixture was incubated at 37°C for 3 hours in the presence of 5% CO₂. After the sample solution was
20 removed, the cell surface was washed with TDMEM, 2 ml per well of DMEM containing 10% fetal calf albumin was added, and the incubation was carried out at 37°C for 2 days in the presence of 5% CO₂.

 To 10 ml of a 50 mM phosphate buffer solution (pH 7.4)
25 containing 2% bovine fibrinogen (Miles Inc.), 0.5% agarose, and 1 mM calcium chloride were added 10 units of human thrombin (Mochida Pharmaceutical Co., Ltd.) and the resulting mixture was solidified in a plate of 9 cm in diameter to prepare a fibrin plate. Ten microliters of the
30 culture supernatant of the tansfected COS7 cells were

spotted on the fibrin plate, which was incubated at 37°C for 15 hours. In the case in which a clear circle appears on the fibrin plate, it is judged that the cDNA fragment codes for the amino acid sequence functioning as a secretory signal sequence. On the other hand, in case in which a clear circle is not formed, the cells were washed well, then the fibrin sheet was placed on the cells, and incubation was carried out at 37°C for 15 hours. In case in which a clear portion is formed on the fibrin sheet, it indicates that the urokinase activity was expressed on the cell surface. In other words, the cDNA fragment is judged to code for the transmembrane domains.

(5) Protein Synthesis by In Vitro Translation

The plasmid vector bearing the cDNA of the present invention was used for in vitro transcription/translation with a T_NT rabbit reticulocyte lysate kit (Promega). In this case, [³⁵S]methionine was added to label the expression product with a radioisotope. Each of the reactions was carried out according to the protocols attached to the kit. Two micrograms of the plasmid was reacted at 30°C for 90 minutes in a total 25 µl volume of the reaction solution containing 12.5 µl of T_NT rabbit reticulocyte lysate, 0.5 µl of a buffer solution (attached to kit), 2 µl of an amino acid mixture (methionine-free), 2 µl of [³⁵S]methionine (Amersham) (0.37 MBq/µl), 0.5 µl of T7RNA polymerase, and 20 U of RNasin. To 3 µl of the resulting reaction solution was added 2 µl of the SDS sampling buffer (125 mM Tris-hydrochloric acid buffer, pH 6.8, 120 mM 2-mercaptoethanol, 2% SDS solution, 0.025% bromophenol blue, and 20% glycerol) and the resulting

mixture was heated at 95°C for 3 minutes and then subjected to SDS-polyacrylamide gel electrophoresis. The molecular weight of the translation product was determined by carrying out the autoradiograph.

5 (6) Expression by COS7

Escherichia coli bearing the expression vector of the protein of the present invention was infected with helper phage M13K07 and single-stranded phage particles were obtained by the above-mentioned procedure. The thus-
10 obtained phage was used for introducing each expression vector in the culture cells originating from the simian kidney, COS7. After incubation at 37°C for 2 days in the presence of 5% CO₂, the incubation was continued for one hour in the culture medium containing [³⁵S]cystine or
15 [³⁵S]methionine. Collection and dissolution of the cells, followed by subjecting to SDS-PAGE, allowed to observe the presence of a band corresponding to the expression product of each protein, which did not exist in the COS7 cells. For instance, HP01207 produced a band of 25 kDa in the
20 membrane fraction.

(7) Clone Examples

<HP01207> (Sequence Nos. 1, 4, and 7)

Determination of the whole base sequence of the cDNA insert of clone HP01207 obtained from cDNA libraries of
25 human stomach cancer revealed the structure consisting of a 100-bp 5'-nontranslation region, an 810-bp ORF, and a 2028-bp 3'-nontranslation region. The ORF codes for a protein consisting of 269 amino acid residues and there existed seven transmembrane domains. Figure 1 depicts the
30 hydrophobicity/hydrophilicity profile, obtained by the

Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a smear translation product of a high molecular weight.

The search of the protein data base by using the amino acid sequence of the present protein revealed that the protein was analogous to the mouse Surf-4 protein (PIR Accession No. A34727). Table 2 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the mouse Surf-4 protein (MM). Therein, the marks of * and . represent an amino acid residue identical with the protein of the present invention and an amino acid residue analogous to the protein of the present invention, respectively. The both proteins possessed a homology of 99.3% in the entire region.

Table 2

	HS	MGQNDLMGTAEDFADQFLRVTKQYLPHVARLCLISTFLEDGIRMWFQWSEQRDYIDTTWN
		*****.
20	MM	MGQNDLMGTAEDFADQFLRVTKQYLPHVARLCLISTFLEDGIRMWFQWSEQRDYIDTTWS
	HS	CGYLLASSFVFLNLLGQLTGCVLVLSRNFVQYACFGLFGIIALQTIAYSILWDLKFLMRN

	MM	CGYLLASSFVFLNLLGQLTGCVLVLSRNFVQYACFGLFGIIALQTIAYSILWDLKFLMRN
	HS	LALGGGLLLLLAESRSEGKSMFAGVPTMRESSPKQYMLGGRVLLVLMFMTLLHFDASFF
25		*****
	MM	LALGGGLLLLLAESRSEGKSMFAGVPTMRESSPKQYMLGGRVLLVLMFMTLLHFDASFF
	HS	SIVQNIVGTALMILVAIGFKTKLAALTLVVWLFAINVYFNAFWTIPVYKPMHDFLKYDF
		**.*
	MM	SIIQNIVGTALMILVAIGFKTKLAALTLVVWLFAINVYFNAFWTIPVYKPMHDFLKYDF
30	HS	QTMSVIGGLLLVVALGPGGVSMDEKKKEW

	MM	QTMSVIGGLLLVVALGPGGVSMDEKKKEW

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of a base sequence that exhibited an analogy of 98.6% with a 762-bp part from position 122 up to position 883 (GenBank Accession No. Y14820), which codes for the fragment of the present protein.

The mouse Surf-4 protein is one of proteins which are encoded in the mouse surfeit locus and has been considered to a housekeeping protein that is essential to the survival of cells [Huxley, C. et al., Mol. Cell. Biol. 10: 605-614 (1990)].

<HP01862> (Sequence Nos. 2, 5 and 9)

Determination of the whole base sequence of the cDNA insert of clone HP01862 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of an 80-bp 5'-nontranslation region, a 936-bp ORF, and a 1274-bp 3'-nontranslation region. The ORF codes for a protein consisting of 311 amino acid residues and there existed seven transmembrane domains. Figure 2 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a smear translation product of a high molecular weight.

The search of the protein data base using the amino acid sequence of the present protein has revealed the presence of sequences that were analogous to the rat NMDA receptor glutamate-binding subunit (GenBank Accession No. S19586). Table 3 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the rat NMDA receptor glutamate-binding subunit

5

10

15

30

found to contain the initiation codon.

The rat NMDA receptor glutamate-binding subunit is one of subunits of an NMDA receptor complex which exist specifically in the brain [Kumar, K. N. et al., Nature 354: 70-73 (1991)]. The protein of the present invention has seven transmembrane domains characteristic to channels and transporters and thereby is considered to play an important role as a channel and a transporter.

<HP10493> (Sequence Nos. 3, 6 and 11)

Determination of the whole base sequence of the cDNA insert of clone HP10493 obtained from cDNA libraries of the human lymphoma U937 revealed the structure consisting of a 123-bp 5'-nontranslation region, a 1152-bp ORF, and a 2430-bp 3'-nontranslation region. The ORF codes for a protein consisting of 383 amino acid residues and there existed one transmembrane domain at the N-terminus. Figure 3 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. Introduction of an expression vector, wherein the HindIII-AccI fragment containing a cDNA portion coding for the N-terminal 44 amino acid residues of the present protein was inserted into the HindIII-PmaCI site of pSSD3, into the COS7 cells revealed the urokinase activity on the cell surface to indicate that the present protein is the type-II membrane protein. In vitro translation resulted in formation of a translation product of 43 kDa that was almost consistent with the molecular weight of 43,001 predicted from the ORF.

The search of the protein data base using the amino acid sequence of the present protein has not revealed the presence of any known protein having an analogy. The search

of the motif sequences has revealed a high probability that histidine at position 175 is an active site of the trypsin-type serine protease. Accordingly, the present protein is likely to be a membrane-type protease. Also, the GenBank
5 using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. R81003) in EST, but many sequences were not distinct and the same ORF as that in the present cDNA was not found.

INDUSTRIAL APPLICABILITY

The present invention provides human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic
15 cells expressing said cDNAs. All of the proteins of the present invention exist in the cell membrane, so that they are considered to be proteins controlling the proliferation and the differentiation of the cells. Accordingly, the proteins of the present invention can be employed as
20 pharmaceuticals such as carcinostatic agents relating to the control of the proliferation and the differentiation of the cells or as antigens for preparing antibodies against said proteins. The cDNAs of the present invention can be utilized as probes for the gene diagnosis and gene sources
25 for the gene therapy. Furthermore, the cDNAs can be utilized for large-scale expression of said proteins. Cells, wherein these membrane protein genes are introduced and membrane proteins are expressed in large amounts, can be utilized for detection of the corresponding ligands,
30 screening of novel low-molecular pharmaceuticals, and so on.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are

5 transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding
10 sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods
15 include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that has been separated from the adjacent coding
20 sequences, if any, present in the genome of the organism from which the gene was isolated.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided.
25 The desired change in gene expression can be achieved through the use of antisense polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, Trends Pharmacol. Sci. 15(7): 250-254; Lavarosky et al., 1997, Biochem. Mol. Med. 62(1):
30 11-22; and Hampel, 1998, Prog. Nucleic Acid Res. Mol. Biol.

58: 1-39; all of which are incorporated by reference herein). Transgenic animals that have multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, Bioessays 14(9): 629-633; Zwaal et al., 1993, Proc. Natl. Acad. Sci. USA 90(16): 7431-7435; Clark et al., 1994, Proc. Natl. Acad. Sci. USA 91(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination, preferably detected by positive/negative genetic selection strategies (Mansour et al., 1988, Nature 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms

are useful for the development of non-human models for the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms part or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information.

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity;

most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

Species homologs of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide, as determined by those of skill in the art. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous, or related to that encoded by the polynucleotides.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides capable of hybridizing under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for

example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Table

5

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) [†]	Hybridization Temperature and Buffer [†]	Wash Temperature and Buffer [†]
A	DNA : DNA	≥50	65°C; 1×SSC -or- 42°C; 1×SSC, 50% formamide	65°C; 0.3×SSC
B	DNA : DNA	<50	T _B *; 1×SSC	T _B *; 1×SSC
C	DNA : RNA	≥50	67°C; 1×SSC -or- 45°C; 1×SSC, 50% formamide	67°C; 0.3×SSC
D	DNA : RNA	<50	T _D *; 1×SSC	T _D *; 1×SSC
E	RNA : RNA	≥50	70°C; 1×SSC -or- 50°C; 1×SSC, 50% formamide	70°C; 0.3×SSC
F	RNA : RNA	<50	T _F *; 1×SSC	T _F *; 1×SSC
G	DNA : DNA	≥50	65°C; 4×SSC -or- 42°C; 4×SSC, 50% formamide	65°C; 1×SSC
H	DNA : DNA	<50	T _H *; 4×SSC	T _H *; 4×SSC
I	DNA : RNA	≥50	67°C; 4×SSC -or- 45°C; 4×SSC, 50% formamide	67°C; 1×SSC
J	DNA : RNA	<50	T _J *; 4×SSC	T _J *; 4×SSC
K	RNA : RNA	≥50	70°C; 4×SSC -or- 50°C; 4×SSC, 50% formamide	67°C; 1×SSC
L	RNA : RNA	<50	T _L *; 2×SSC	T _L *; 2×SSC
M	DNA : DNA	≥50	50°C; 4×SSC -or- 40°C; 6×SSC, 50% formamide	50°C; 2×SSC
N	DNA : DNA	<50	T _N *; 6×SSC	T _N *; 6×SSC
O	DNA : RNA	≥50	55°C; 4×SSC -or- 42°C; 6×SSC, 50% formamide	55°C; 2×SSC
P	DNA : RNA	<50	T _P *; 6×SSC	T _P *; 6×SSC
Q	RNA : RNA	≥50	60°C; 4×SSC -or- 45°C; 6×SSC, 50% formamide	60°C; 2×SSC
R	RNA : RNA	<50	T _R *; 4×SSC	T _R *; 4×SSC

‡ : The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

† : SSPE (1×SSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH7.4) can be substituted for SSC (1×SSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

*T_B - T_R : The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, T_m(°C)=2(#of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T_m(°C)=81.5 + 16.6(log₁₀[Na⁺]) + 0.41 (%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium ions in the hybridization buffer ([Na⁺] for 1×SSC=0.165M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning: A Laboratory

Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and Current Protocols in Molecular Biology, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc.,

sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present

invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

CLAIMS

1. A protein comprising any one of the amino acid sequences represented by Sequence Nos. 1 to 3.

5 2. A DNA coding for the protein according to Claim 1.

3. A cDNA comprising any one of the base sequences represented by Sequence Nos. 4 to 6.

10 4. The cDNA according to Claim 3 comprising any one of the base sequences represented by Sequence Nos. 7, 9 and 11.

5. An expression vector capable of expressing the DNA according to any one of Claims 2 to 4 by in vitro translation or in eucaryotic cells.

15 6. A transformation eucaryotic cell capable of expressing the DNA according to any one of Claims 2 to 4 and of producing the protein according to Claim 1.




Fig.1




Fig.2

0954933.082100




Fig.3

0954933.082100

**DECLARATION, PETITION AND POWER OF ATTORNEY
FOR PATENT APPLICATION**

(Check one):

- ☐ Declaration Submitted with Initial Filing
☒ Declaration Submitted after Initial Filing

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**HUMAN PROTEINS HAVING TRANSMEMBRANE DOMAINS AND DNAs
ENCODING THESE PROTEINS**

the specification of which (check one):

☐ is attached hereto.

OR

☒ was filed on **22 May 2000** as U.S. National Application Serial No. 09/554,933
(U.S. National Filing of PCT/JP98/05238 filed on 20 November 1998).

☐ and was amended by PCT Article 19 Amendment on _____
(if applicable),

☐ and was amended by PCT Article 34 Amendment on _____
(if applicable).

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby state that I have reviewed and understood the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

(Check one):

- 1) FOREIGN PRIORITY CLAIM:** I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate or §365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or any PCT international application having a filing date before that of the application on which priority is claimed.

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto.

Provisional Application Number(s)	Filing Date (dd/mm/yyyy)

- 3) **U.S./PCT PRIORITY CLAIM:** I hereby claim the benefit under Title 35, United States Code, §120 of any United States application or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

POWER OF ATTORNEY:

As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

James E. Cockfield Reg. No. 19,162
Thomas V. Smurzynski Reg. No. 24,798
Ralph A. Loren Reg. No. 29,325
Giulio A. DeConti, Jr. Reg. No. 31,503
Ann Lamport Hammitte Reg. No. 34,858
Elizabeth A. Hanley Reg. No. 33,505
Amy E. Mandragouras Reg. No. 36,207
Anthony A. Laurentano Reg. No. 38,220
Jane E. Remillard Reg. No. 38,872
Jeremiah Lynch Reg. No. 17,425
Kevin J. Canning Reg. No. 35,470
David A. Lane, Jr. Reg. No. 39,261
Jeanne M. DiGiorgio Reg. No. 41,710

Megan E. Williams Reg. No. 43,270
Nicholas P. Triano III Reg. No. 36,397
Peter C. Lauro Reg. No. 32,360
DeAnn F. Smith Reg. No. 36,683
William D. DeVaul Reg. No. 42,483
David J. Rikkers Reg. No. 43,882
Chi Suk Kim Reg. No. 42,728
Maria Laccotripe Zacharakis Limited Recognition
Under 37 C.F.R. § 10.9(b)
Debra J. Milasincic Reg. No. P46,931
David R. Burns Reg. No. P46,590

of LAHIVE & COCKFIELD, LLP, 28 State Street, Boston, Massachusetts 02109, United States of America, and

Ellen J. Kapinos Reg. No. 32,245 Elizabeth Hurley Reg. No. 41,859
Barbara A. Gyure Reg. No. 34,614
M. Kymne Hehman Reg. No. 39,206

of GENETICS INSTITUTE, INC., 87 CambridgePark Drive, Cambridge, Massachusetts 02140, United States of America,

Egon E. Berg Reg. No. 21,117 Elizabeth M. Barnhard Reg. No. 31,088
Gale F. Matthews Reg. No. 32,269 Alan M. Gordon Reg. No. 30,637
Darryl L. Webster Reg. No. 34,276

of GENETICS INSTITUTE, INC., One Campus Drive, Parsippany, New Jersey 07054, United States of America, and

Rebecca R. Barrett Reg. No. 35,152 Steven R. Eck Reg. No. 36,126
Arnold S. Milowsky Reg. No. 35,288 Michael R. Nagy Reg. No. 33,432
George Tarnowski Reg. No. 27,472 Arthur G. Seifert Reg. No. 28,040

of GENETICS INSTITUTE, INC., P.O. Box 8299, Philadelphia, Pennsylvania 19101, United States of America.

007680-2264560

Send Correspondence to:

Amy E. Mandragouras, Lahive & Cockfield, LLP, 28 State Street, Boston, Massachusetts 02109, United States of America

Direct Telephone Calls to: (name and telephone number)

Peter C. Lauro, (617) 227-7400, also at Lahive & Cockfield, LLP, 28 State Street, Boston, Massachusetts 02109, United States of America

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Seishi KATO	
Inventor's signature <i>Seishi Kato</i>	Date 10 August 2000
Residence 3-46-50, Wakamatsu, Sagamihara-shi, Kanagawa 229-0014, Japan	
Citizenship Japan	
Post Office Address (if different)	

Full name of sole or first inventor Tomoko KIMURA	
Inventor's signature <i>Tomoko Kimura</i>	Date 10 August 2000
Residence 715, 2-9-1, Kohoku, Tsuchiura-shi, Ibaraki 300-0032, Japan	
Citizenship Japan	
Post Office Address (if different)	

34

526 Rec'd PCT/PTO 22 MAY 2000

28 State Street
Boston, MA 02109
Tel. (617) 227-7400
Dated: **22 May 2000**

SEQUENCE LISTING

<110> Sagami Chemical Research Center

5 <120> Human Proteins Having Transmembrane Domains and DNAs Encoding these Proteins

<130> 660857

<140>

<141>

10

<150> Japan 9-323129

<151> 1997-11-25

<160> 12

15

<170> Windows 95 (Word 98)

<210> 1

<211> 269

20

<212> PRT

<213> Homo sapiens

<400> 1

Met Gly Gln Asn Asp Leu Met Gly Thr Ala Glu Asp Phe Ala Asp Gln

25

1 5 10 15

Phe Leu Arg Val Thr Lys Gln Tyr Leu Pro His Val Ala Arg Leu Cys

20 25 30

Leu Ile Ser Thr Phe Leu Glu Asp Gly Ile Arg Met Trp Phe Gln Trp
 35 40 45
 Ser Glu Gln Arg Asp Tyr Ile Asp Thr Thr Trp Asn Cys Gly Tyr Leu
 50 55 60
 5 Leu Ala Ser Ser Phe Val Phe Leu Asn Leu Leu Gly Gln Leu Thr Gly
 65 70 75 80
 Cys Val Leu Val Leu Ser Arg Asn Phe Val Gln Tyr Ala Cys Phe Gly
 85 90 95
 Leu Phe Gly Ile Ile Ala Leu Gln Thr Ile Ala Tyr Ser Ile Leu Trp
 10 100 105 110
 Asp Leu Lys Phe Leu Met Arg Asn Leu Ala Leu Gly Gly Gly Leu Leu
 115 120 125
 Leu Leu Leu Ala Glu Ser Arg Ser Glu Gly Lys Ser Met Phe Ala Gly
 130 135 140
 15 Val Pro Thr Met Arg Glu Ser Ser Pro Lys Gln Tyr Met Gln Leu Gly
 145 150 155 160
 Gly Arg Val Leu Leu Val Leu Met Phe Met Thr Leu Leu His Phe Asp
 165 170 175
 Ala Ser Phe Phe Ser Ile Val Gln Asn Ile Val Gly Thr Ala Leu Met
 20 180 185 190
 Ile Leu Val Ala Ile Gly Phe Lys Thr Lys Leu Ala Ala Leu Thr Leu
 195 200 205
 Val Val Trp Leu Phe Ala Ile Asn Val Tyr Phe Asn Ala Phe Trp Thr
 210 215 220
 25 Ile Pro Val Tyr Lys Pro Met His Asp Phe Leu Lys Tyr Asp Phe Phe
 225 230 235 240
 Gln Thr Met Ser Val Ile Gly Gly Leu Leu Leu Val Val Ala Leu Gly

245 250 255
Pro Gly Gly Val Ser Met Asp Glu Lys Lys Lys Glu Trp

260 265

5 <210> 2

<211> 311

<212> PRT

<213> Homo sapiens

10 <400> 2

Met Ser Asn Pro Ser Ala Pro Pro Pro Tyr Glu Asp Arg Asn Pro Leu

1 5 10 15

Tyr Pro Gly Pro Leu Pro Pro Gly Gly Tyr Gly Gln Pro Ser Val Leu

20 25 30

15 Pro Gly Gly Tyr Pro Ala Tyr Pro Gly Tyr Pro Gln Pro Gly Tyr Gly

35 40 45

His Pro Ala Gly Tyr Pro Gln Pro Met Pro Pro Thr His Pro Met Pro

50 55 60

Met Asn Tyr Gly Pro Gly His Gly Tyr Asp Gly Glu Glu Arg Ala Val

20 65 70 75 80

Ser Asp Ser Phe Gly Pro Gly Glu Trp Asp Asp Arg Lys Val Arg His

85 90 95

Thr Phe Ile Arg Lys Val Tyr Ser Ile Ile Ser Val Gln Leu Leu Ile

100 105 110

25 Thr Val Ala Ile Ile Ala Ile Phe Thr Phe Val Glu Pro Val Ser Ala

115 120 125

Phe Val Arg Arg Asn Val Ala Val Tyr Tyr Val Ser Tyr Ala Val Phe

130 135 140
 Val Val Thr Tyr Leu Ile Leu Ala Cys Cys Gln Gly Pro Arg Arg Arg
 145 150 155 160
 Phe Pro Trp Asn Ile Ile Leu Leu Thr Leu Phe Thr Phe Ala Met Gly
 5 165 170 175
 Phe Met Thr Gly Thr Ile Ser Ser Met Tyr Gln Thr Lys Ala Val Ile
 180 185 190
 Ile Ala Met Ile Ile Thr Ala Val Val Ser Ile Ser Val Thr Ile Phe
 195 200 205
 10 Cys Phe Gln Thr Lys Val Asp Phe Thr Ser Cys Thr Gly Leu Phe Cys
 210 215 220
 Val Leu Gly Ile Val Leu Leu Val Thr Gly Ile Val Thr Ser Ile Val
 225 230 235 240
 Leu Tyr Phe Gln Tyr Val Tyr Trp Leu His Met Leu Tyr Ala Ala Leu
 15 245 250 255
 Gly Ala Ile Cys Phe Thr Leu Phe Leu Ala Tyr Asp Thr Gln Leu Val
 260 265 270
 Leu Gly Asn Arg Lys His Thr Ile Ser Pro Glu Asp Tyr Ile Thr Gly
 275 280 285
 20 Ala Leu Gln Ile Tyr Thr Asp Ile Ile Tyr Ile Phe Thr Phe Val Leu
 290 295 300
 Gln Leu Met Gly Asp Arg Asn
 305 310
 25 <210> 3
 <211> 383
 <212> PRT

<213> Homo sapiens

<400> 3

Met Ala Gly Ile Pro Gly Leu Leu Phe Leu Leu Phe Phe Leu Leu Cys

5 1 5 10 15

Ala Val Gly Gln Val Ser Pro Tyr Ser Ala Pro Trp Lys Pro Thr Trp

20 25 30

Pro Ala Tyr Arg Leu Pro Val Val Leu Pro Gln Ser Thr Leu Asn Leu

35 40 45

10 Ala Lys Pro Asp Phe Gly Ala Glu Ala Lys Leu Glu Val Ser Ser Ser

50 55 60

Cys Gly Pro Gln Cys His Lys Gly Thr Pro Leu Pro Thr Tyr Glu Glu

65 70 75 80

Ala Lys Gln Tyr Leu Ser Tyr Glu Thr Leu Tyr Ala Asn Gly Ser Arg

15 85 90 95

Thr Glu Thr Gln Val Gly Ile Tyr Ile Leu Ser Ser Ser Gly Asp Gly

100 105 110

Ala Gln His Arg Asp Ser Gly Ser Ser Gly Lys Ser Arg Arg Lys Arg

115 120 125

20 Gln Ile Tyr Gly Tyr Asp Ser Arg Phe Ser Ile Phe Gly Lys Asp Phe

130 135 140

Leu Leu Asn Tyr Pro Phe Ser Thr Ser Val Lys Leu Ser Thr Gly Cys

145 150 155 160

Thr Gly Thr Leu Val Ala Glu Lys His Val Leu Thr Ala Ala His Cys

25 165 170 175

Ile His Asp Gly Lys Thr Tyr Val Lys Gly Thr Gln Lys Leu Arg Val

180 185 190

Gly Phe Leu Lys Pro Lys Phe Lys Asp Gly Gly Arg Gly Ala Asn Asp

195 200 205

Ser Thr Ser Ala Met Pro Glu Gln Met Lys Phe Gln Trp Ile Arg Val

210 215 220

5 Lys Arg Thr His Val Pro Lys Gly Trp Ile Lys Gly Asn Ala Asn Asp

225 230 235 240

Ile Gly Met Asp Tyr Asp Tyr Ala Leu Leu Glu Leu Lys Lys Pro His

245 250 255

Lys Arg Lys Phe Met Lys Ile Gly Val Ser Pro Pro Ala Lys Gln Leu

10 260 265 270

Pro Gly Gly Arg Ile His Phe Ser Gly Tyr Asp Asn Asp Arg Pro Gly

275 280 285

Asn Leu Val Tyr Arg Phe Cys Asp Val Lys Asp Glu Thr Tyr Asp Leu

290 295 300

15 Leu Tyr Gln Gln Cys Asp Ala Gln Pro Gly Ala Ser Gly Ser Gly Val

305 310 315 320

Tyr Val Arg Met Trp Lys Arg Gln Gln Gln Lys Trp Glu Arg Lys Ile

325 330 335

Ile Gly Ile Phe Ser Gly His Gln Trp Val Asp Met Asn Gly Ser Pro

20 340 345 350

Gln Asp Phe Asn Val Ala Val Arg Ile Thr Pro Leu Lys Tyr Ala Gln

355 360 365

Ile Cys Tyr Trp Ile Lys Gly Asn Tyr Leu Asp Cys Arg Glu Gly

370 375 380

25

<210> 4

<211> 807

<213> Homo sapiens

5 atgggcagac acgacctgat gggcacggcc gaggacttcg ccgaccagtt cctccgtgtc 60
acaaagcagt acctgccccac cgtggcgcgcc ctctgtctga tcagcacctt cctggaggac 120
ggcatccgta tgtggttcca gtggagcgag cagcgcgact acatcgacac cacctggaac 180
tgcggctacc tgctggcctc gtccttcgtc ttctcaact tgctgggaca gctgactggc 240
tgcgtcctgg tgttgagcag gaacttcgtg cagtacgcct gcttcgggct ctttggaaac 300
10 atagctctgc agacgattgc ctacagcatt ttatgggact tgaagtttt gatgaggaac 360
ctggccctgg gaggaggcct gttgctgctc ctagcagaat cccgttctga agggaagagc 420
atgtttgcgg gcgtcccccac catgcgtgag agctccccca aacagtacat gcagctcgga 480
ggcagggtct tgctggttct gatgttcatt accctccttc actttgacgc cagcttcttt 540
tctattgtcc agaacatcgt gggcacagct ctgatgattt tagtggccat tggttttaaa 600
15 accaagctgg ctgctttgac tcttgttgtg tggctctttg ccatcaacgt atatttcaac 660
gccttctgga ccattccagt ctacaagccc atgcatgact tcttgaaata cgacttcttc 720
cagaccatgt cggtgattgg gggcttgctc ctggtggtgg ccttggggcc tgggggtgtc 780
tccatggatg agaagaagaa ggagtgg 807

<211> 933

<213> Homo sapiens

atgtccaacc ccagcgcccc accaccatfat gaagaccgca accccctgta cccaggccct 60
ctgccccctg ggggctatgg gcagccatct gtctctgccag gagggtatcc tgccctacct 120

ggctaccgc agcctggcta cggtcaccct gctggctacc cacagcccat gccccccacc 180
 caccgatgc ccatgaacta cggcccaggc catggctatg atggggagga gagagcgggtg 240
 agtgatagct tcgggcctgg agagtgggat gaccggaaag tgcgacacac tttatccga 300
 aaggtttact ccatcatctc cgtgcagctg ctcatcactg tggccatcat tgctatcttc 360
 5 acctttgtgg aacctgtcag cgcctttgtg aggagaaatg tggctgtcta ctacgtgtcc 420
 tatgtgtct tcgtgtcac ctacctgac ctgacctgct gccagggacc cagacgccgt 480
 ttccatgga acatcattct gctgacctt tttactttg ccatgggctt catgacgggc 540
 accattcca gtatgtacca aaccaaagcc gtcacattg caatgatcat cactgcgggtg 600
 gtatccattt cagtcacat ctctgcttt cagaccaagg tggacttcac ctggtgcaca 660
 10 ggctcttct gtgtcctggg aattgtgctc ctggtgactg ggattgtcac tagcattgtg 720
 ctctacttc aatacgtta ctggctccac atgctctatg ctgctctggg ggccatttgt 780
 ttaccctgt tcttggtta cgacacacag ctggtcctgg ggaaccggaa gcacaccatc 840
 agccccgagg actacatcac tggcgccctg cagatttaca cagacatcat ctacatcttc 900
 acctttgtgc tgcagctgat gggggatcgc aat 933
 15
 <210> 6
 <211> 1149
 <212> DNA
 <213> Homo sapiens
 20
 <400> 6
 atggcaggga ttccagggt cctcttctt ctcttcttc tgctctgtgc tgttgggcaa 60
 gtgagccctt acagtcccc ctggaaacct acttggcctg cataccgctt cctgtcgtc 120
 ttgccccagt ctacctcaa ttagccaag ccagacttg gagccgaagc caaattagaa 180
 25 gtatcttctt catgtggacc ccagtgtcat aagggaactc cactgcccac ttacgaagag 240
 gccaaagcaat atctgtctta tgaacgctc tatgccaatg gcagccgcac agagacgcag 300
 gtgggcatct acatctcag cagtagtgga gatggggccc aacaccgaga ctcagggtct 360

tcaggaaagt ctggaaggaa gcggcagatt tatggctatg acagcagggt cagcatttt 420
gggaaggact tctgtctcaa ctaccctttc tcaacatcag tgaagtatc cacgggctgc 480
accggcaccc tgggtggcaga gaagcatgtc ctacacagctg cccactgcac acacgatgga 540
aaaacctatg tgaaaggaac ccagaagctt cgagtgggct tctaaagcc caagtttaaa 600
5 gatgtgtgtc gagggggccaa cgactccact tcagccatgc ccgagcagat gaaatttcag 660
tggatccggg tgaaacgcac ccatgtgccc aagggttga tcaagggcaa tgccaatgac 720
atcggcatgg attatgatta tgccctcctg gaactcaaaa agccccacaa gagaaaattt 780
atgaagattg ggggtagccc tctgtctaag cagctgccag ggggcagaat tcacttctct 840
ggttatgaca atgaccgacc aggcaatttg gtgtatcgt tctgtgacgt caaagacgag 900
10 acctatgact tgctctacca gcaatgcgat gccagccag gggccagcgg gtctggggtc 960
tatgtgagga tgtggaagag acagcagcag aagtgggagc gaaaaattat tggcattttt 1020
tcagggcacc agtgggtgga catgaatgt tccccacagg attcaacgt ggctgtcaga 1080
atcactcctc tcaaatatgc ccagatttc tattggatta aaggaaacta cctggattgt 1140
agggagggg 1149
15
<210> 7
<211> 2938
<212> DNA
<213> Homo sapiens
20
<400> 7
aaaaagggca ctctctgtgg aggccgcagc ggggtgcgggc gccgacgggc gagagccagc 60
gagcgagcga gcgagccgag ccgagcctcc cgccgtgcc atg ggc cag aac gac 115
Met Gly Gln Asn Asp
25 1 5
ctg atg ggc acg gcc gag gac ttc gcc gac cag ttc ctc cgt gtc aca 163
Leu Met Gly Thr Ala Glu Asp Phe Ala Asp Gln Phe Leu Arg Val Thr

10 15 20

aag cag tac ctg ccc cac gtg gcg cgc ctc tgt ctg atc agc acc ttc 211

Lys Gln Tyr Leu Pro His Val Ala Arg Leu Cys Leu Ile Ser Thr Phe

25 30 35

5 ctg gag gac ggc atc cgt atg tgg ttc cag tgg agc gag cag cgc gac 259

Leu Glu Asp Gly Ile Arg Met Trp Phe Gln Trp Ser Glu Gln Arg Asp

40 45 50

tac atc gac acc acc tgg aac tgc ggc tac ctg ctg gcc tgc tcc ttc 307

Tyr Ile Asp Thr Thr Trp Asn Cys Gly Tyr Leu Leu Ala Ser Ser Phe

10 55 60 65

gtc ttc ctc aac ttg ctg gga cag ctg act ggc tgc gtc ctg gtg ttg 355

Val Phe Leu Asn Leu Leu Gly Gln Leu Thr Gly Cys Val Leu Val Leu

70 75 80 85

agc agg aac ttc gtg cag tac gcc tgc ttc ggg ctc ttt gga atc ata 403

15 Ser Arg Asn Phe Val Gln Tyr Ala Cys Phe Gly Leu Phe Gly Ile Ile

90 95 100

gct ctg cag acg att gcc tac agc att tta tgg gac ttg aag ttt ttg 451

Ala Leu Gln Thr Ile Ala Tyr Ser Ile Leu Trp Asp Leu Lys Phe Leu

105 110 115

20 atg agg aac ctg gcc ctg gga gga ggc ctg ttg ctg ctc cta gca gaa 499

Met Arg Asn Leu Ala Leu Gly Gly Gly Leu Leu Leu Leu Leu Ala Glu

120 125 130

tcc cgt tct gaa ggg aag agc atg ttt gcg ggc gtc ccc acc atg cgt 547

Ser Arg Ser Glu Gly Lys Ser Met Phe Ala Gly Val Pro Thr Met Arg

25 135 140 145

gag agc tcc ccc aaa cag tac atg cag ctc gga ggc agg gtc ttg ctg 595

Glu Ser Ser Pro Lys Gln Tyr Met Gln Leu Gly Gly Arg Val Leu Leu

150 155 160 165

gtt ctg atg ttc atg acc ctc ctt cac ttt gac gcc agc ttc ttt tct 643

Val Leu Met Phe Met Thr Leu Leu His Phe Asp Ala Ser Phe Phe Ser

170 175 180

5 att gtc cag aac atc gtg ggc aca gct ctg atg att tta gtg gcc att 691

Ile Val Gln Asn Ile Val Gly Thr Ala Leu Met Ile Leu Val Ala Ile

185 190 195

ggg ttt aaa acc aag ctg gct gct ttg act ctt gtt gtg tgg ctc ttt 739

Gly Phe Lys Thr Lys Leu Ala Ala Leu Thr Leu Val Val Trp Leu Phe

10 200 205 210

gcc atc aac gta tat ttc aac gcc ttc tgg acc att cca gtc tac aag 787

Ala Ile Asn Val Tyr Phe Asn Ala Phe Trp Thr Ile Pro Val Tyr Lys

215 220 225

ccc atg cat gac ttc ctg aaa tac gac ttc ttc cag acc atg tcg gtg 835

15 Pro Met His Asp Phe Leu Lys Tyr Asp Phe Phe Gln Thr Met Ser Val

230 235 240 245

att ggg ggc ttg ctc ctg gtg gtg gcc ctg ggc cct ggg ggt gtc tcc 883

Ile Gly Gly Leu Leu Leu Val Val Ala Leu Gly Pro Gly Gly Val Ser

250 255 260

20 atg gat gag aag aag aag gag tgg taa cagtcacaga tcctacctg 930

Met Asp Glu Lys Lys Lys Glu Trp

265

cctggctaag acccgtagcc gtcaaggact gggtcggggg ggaatcaaca aaactgccag 990

cttttatgta tctcttccc ttccccccc ttggtaaagg cacagatggt ttgagaactt 1050

25 tatttgacaga gacacctgag aatcgatggc tcagtctgct ctggagccac agtctggcgt 1110

ctgacccttc agtgcaggcc agcctggcag ctggaagcct cccccacgcc gaggctttgg 1170

agtgaacagc ccgcttggtc gtggcatctc agtctctatt ttgagtttt ttgtgggggt 1230

acaggagggg gccttcaagc tgtactgtga gcagacgcat tggattatc attcaaagca 1290
gtctccctct tatttgtaag ttacatttt tagcggaac tactaaatta tttgggtgg 1350
ttcagccaaa cctcaaaaca gtaatctcc ctggtttaa atcacaccag tggtttgat 1410
gttgtttctg ccccgattg tttttatag gaatactgaa aacatttagg gacacccaaa 1470
5 gaatgatgca gtattaaagg ggtggtagaa gctgctgtt atgataaaag tcatcggtca 1530
gaaaatcagc ttggattgt gccaagtgt ttattgggta acaccctggg agttttagta 1590
gcttgaggca aggtggaggg gcaagaagtc ctggggaag ctgctggtct gggtgctgct 1650
ggcctccaag ctggcagtgg gaagggtag tgagaccaca caggggtagc cccagcagca 1710
gcaccctgca agccagcctg gccagctgct cagaccagct tgcagagccg cagccgctgt 1770
10 gggcaggggg tgtggcagga gctcccagca ctggagacc accgactcaa cccagttacc 1830
tcacatgggg cttttctga gcaaggtctc gaaagcgcag gccgccctgg ctgagcagca 1890
ccgcccttc ccagctgcac tcgccctgtg gacagccccg acacaccact ttctgaggc 1950
tgtcgtcac tcagattgtc cgtttgctat gccgaatgca gccaaaattc cttttacaa 2010
tttgtgatgc ctaccgatt tgattctaat cctgtattta aagttttcta aactgcctt 2070
15 atactgtgtt tctcttttg ggggagctta actgcttgtt gctccctgtc gtctgcacca 2130
tagtaaatgc cacaagggtg gtcgaacacc tctctggccc ctagacctat ctggggacag 2190
gttggtcag cctgtctcca gggtgctgc ggcccagccc cgagcctgcc tccctcttg 2250
cctctcatcc attggctctg cagggcaggg gtgaggcagg ttctgctca taagtgttt 2310
tggaagtcac ctacctttt aacacagccg aactagtccc aacgcgttg caaatattcc 2370
20 cctggtagcc tacttctta ccccggaata ttgtaagat cgatcaatgg ctcaggaca 2430
tgggttctct tctcctgtga tcattcaagt gtcactgca tgaagactgg cttgtctcag 2490
tgtttcaacc tcaccagggc tgtctcttg tccacacctc gctccctgtt agtgccgtat 2550
gacagcccc atcaaatgac ctggccaag tcacggttc tctgtgtgca aggttggtg 2610
gtgtattgtt ggaaagtagg gtggacaaa ggaggccacg tgagcagtca gcaccagttc 2670
25 tgcaccagca gcgcctccgt ctagtgggt gttctgttt ctctggccc tgggtgggt 2730
agggcctgat tcgggaagat gccttgcag ggaggggagg ataagtggga tctaccaatt 2790
gattctggca aaacaatttc taagatttt ttgctttatg tgggaacag atctaaatct 2850

catttatgc tgtatttat atcttagttg tgtttgaaaa cgttttgatt ttggaaaca 2910

catcaaaata aataatggcg ttgttgt 2938

<210> 8

5 <211> 269

<212> PRT

<213> Homo sapiens

<400> 8

10

Met Gly Gln Asn Asp

1 5

Leu Met Gly Thr Ala Glu Asp Phe Ala Asp Gln Phe Leu Arg Val Thr

10 15 20

Lys Gln Tyr Leu Pro His Val Ala Arg Leu Cys Leu Ile Ser Thr Phe

15

25 30 35

Leu Glu Asp Gly Ile Arg Met Trp Phe Gln Trp Ser Glu Gln Arg Asp

40 45 50

Tyr Ile Asp Thr Thr Trp Asn Cys Gly Tyr Leu Leu Ala Ser Ser Phe

55 60 65

20

Val Phe Leu Asn Leu Leu Gly Gln Leu Thr Gly Cys Val Leu Val Leu

70 75 80 85

Ser Arg Asn Phe Val Gln Tyr Ala Cys Phe Gly Leu Phe Gly Ile Ile

90 95 100

Ala Leu Gln Thr Ile Ala Tyr Ser Ile Leu Trp Asp Leu Lys Phe Leu

25

105 110 115

Met Arg Asn Leu Ala Leu Gly Gly Gly Leu Leu Leu Leu Ala Glu

120 125 130

Met Gly Gln Asn Asp

1 5 10
gac cgc aac ccc ctg tac cca ggc cct ctg ccc cct ggg ggc tat ggg 161
Asp Arg Asn Pro Leu Tyr Pro Gly Pro Leu Pro Pro Gly Gly Tyr Gly
5 15 20 25
cag cca tct gtc ctg cca gga ggg tat cct gcc tac cct ggc tac ccg 209
Gln Pro Ser Val Leu Pro Gly Gly Tyr Pro Ala Tyr Pro Gly Tyr Pro
 30 35 40
cag cct ggc tac ggt cac cct gct ggc tac cca cag ccc atg ccc ccc 257
10 Gln Pro Gly Tyr Gly His Pro Ala Gly Tyr Pro Gln Pro Met Pro Pro
 45 50 55
acc cac ccg atg ccc atg aac tac ggc cca ggc cat ggc tat gat ggg 305
Thr His Pro Met Pro Met Asn Tyr Gly Pro Gly His Gly Tyr Asp Gly
60 65 70 75
15 gag gag aga gcg gtg agt gat agc ttc ggg cct gga gag tgg gat gac 353
Glu Glu Arg Ala Val Ser Asp Ser Phe Gly Pro Gly Glu Trp Asp Asp
 80 85 90
cgg aaa gtg cga cac act ttt atc cga aag gtt tac tcc atc atc tcc 401
Arg Lys Val Arg His Thr Phe Ile Arg Lys Val Tyr Ser Ile Ile Ser
20 95 100 105
gtg cag ctg ctc atc act gtg gcc atc att gct atc ttc acc ttt gtg 449
Val Gln Leu Leu Ile Thr Val Ala Ile Ile Ala Ile Phe Thr Phe Val
 110 115 120
gaa cct gtc agc gcc ttt gtg agg aga aat gtg gct gtc tac tac gtg 497
25 Glu Pro Val Ser Ala Phe Val Arg Arg Asn Val Ala Val Tyr Tyr Val
 125 130 135
tcc tat gct gtc ttc gtt gtc acc tac ctg atc ctt gcc tgc tgc cag 545

Asp Tyr Ile Thr Gly Ala Leu Gln Ile Tyr Thr Asp Ile Ile Tyr Ile

285 290 295

ttc acc ttt gtg ctg cag ctg atg ggg gat cgc aat taaggag 1020

Phe Thr Phe Val Leu Gln Leu Met Gly Asp Arg Asn

5 300 305 310

caagccccc tttcacccg atcctgggct ctcccttcca agctagaggg ctgggccta 1080

tgactgtgt ctgggcttta ggcccttct ctcccttctg agtaacatgc ccagtttct 1140

ttctgtcctg gagacaggtg gcctctctgg ctatggatgt gtgggtactt ggtggggacg 1200

gaggagctag ggactaactg ttgtcttgg tgggcttggc agggactagg ctgaagatgt 1260

10 gtcttctccc cgccacctac tgtatgacac cacattcttc ctaacagctg gggttgtgag 1320

gaatatgaaa agagcctatt ctagtagctag aagggaatat gaaaggtaga agtgacttca 1380

aggtcacgag gttccctccc cacctctgtc acaggcttct tgactacgta gttggagcta 1440

ttcttcccc cagcaaagcc agagagcttt gtcccccggc tcttgacac ataggccatt 1500

atcctgtatt cctttggctt ggcatctttt agctcaggaa ggtagaagag atctgtgccc 1560

15 atgggtctcc ttgttcaat ccctcttctt ttcagtgaca tatgtattgt ttatctgggt 1620

tagggatggg ggacagataa tagaacgagc aaagtaacct atacaggcca gcatggaaca 1680

gcatctcccc tgggcttct cctggcttct gacgtataa gacagagcag gccacatgtg 1740

gccatctgct cccattctt gaaagctgct ggggcctcct tgcaggcttc tggatctctg 1800

gtcagagtga actcttgctt cctgtattca ggcagctcag agcagaaaagt aaggggcaga 1860

20 gtcatacgtg tggccaggaa gtagccaggg tgaagagaga ctcggtgcgg gcaggagaga 1920

tgccctggggg tcctcacct ggctagggag ataccgaagc ctactgtgtt actgaagact 1980

tctgggttct ttcttctgc taaccaggg aggtctctaa gaggaagggt acttctctct 2040

gtttgtctta agttgcactg ggggatttct gacttgaggc ccatctctcc agccagccac 2100

tgccctcttt gtaataataa gtgccttgag ctggaatggg gaagggggac aagggtcagt 2160

25 ctgtcgggtg ggggcagaaa tcaatcagc ccaaggatat agttaggatt aattacttaa 2220

tagagaaatc ctaactatat cacacaaagg gatacaacta taaatgtaat aaaatttatg 2280

tctagaagtt 2290

<210> 10

<211> 311

<212> PRT

5 <213> Homo sapiens

<400> 10

Met Ser Asn Pro Ser Ala Pro Pro Pro Tyr Glu

1 5 10

10 Asp Arg Asn Pro Leu Tyr Pro Gly Pro Leu Pro Pro Gly Gly Tyr Gly

15 20 25

Gln Pro Ser Val Leu Pro Gly Gly Tyr Pro Ala Tyr Pro Gly Tyr Pro

30 35 40

Gln Pro Gly Tyr Gly His Pro Ala Gly Tyr Pro Gln Pro Met Pro Pro

15 45 50 55

Thr His Pro Met Pro Met Asn Tyr Gly Pro Gly His Gly Tyr Asp Gly

60 65 70 75

Glu Glu Arg Ala Val Ser Asp Ser Phe Gly Pro Gly Glu Trp Asp Asp

80 85 90

20 Arg Lys Val Arg His Thr Phe Ile Arg Lys Val Tyr Ser Ile Ile Ser

95 100 105

Val Gln Leu Leu Ile Thr Val Ala Ile Ile Ala Ile Phe Thr Phe Val

110 115 120

Glu Pro Val Ser Ala Phe Val Arg Arg Asn Val Ala Val Tyr Tyr Val

25 125 130 135

Ser Tyr Ala Val Phe Val Val Thr Tyr Leu Ile Leu Ala Cys Cys Gln

140 145 150 155

Gly Pro Arg Arg Arg Phe Pro Trp Asn Ile Ile Leu Leu Thr Leu Phe

160 165 170

Thr Phe Ala Met Gly Phe Met Thr Gly Thr Ile Ser Ser Met Tyr Gln

175 180 185

5 Thr Lys Ala Val Ile Ile Ala Met Ile Ile Thr Ala Val Val Ser Ile

190 195 200

Ser Val Thr Ile Phe Cys Phe Gln Thr Lys Val Asp Phe Thr Ser Cys

205 210 215

Thr Gly Leu Phe Cys Val Leu Gly Ile Val Leu Leu Val Thr Gly Ile

10 220 225 230 235

Val Thr Ser Ile Val Leu Tyr Phe Gln Tyr Val Tyr Trp Leu His Met

240 245 250

Leu Tyr Ala Ala Leu Gly Ala Ile Cys Phe Thr Leu Phe Leu Ala Tyr

255 260 265

15 Asp Thr Gln Leu Val Leu Gly Asn Arg Lys His Thr Ile Ser Pro Glu

270 275 280

Asp Tyr Ile Thr Gly Ala Leu Gln Ile Tyr Thr Asp Ile Ile Tyr Ile

285 290 295

Phe Thr Phe Val Leu Gln Leu Met Gly Asp Arg Asn

20 300 305 310

<210> 11

<211> 3705

<212> DNA

25 <213> Homo sapiens

<400> 11

actctcggt gtgcggcggg gcaggcatgg gagccgcgcg ctcttcccg gcgcccacac 60
ctgtctgagc ggcgacagcga gccgcggccc gggcgggctg ctgcggcggg aacagtgtc 120
ggc atg gca ggg att cca ggg ctc ctc ttc ctt ctc ttc ttt ctg ctc 168
Met Ala Gly Ile Pro Gly Leu Leu Phe Leu Leu Phe Phe Leu Leu
5 1 5 10 15
tgt gct gtt ggg caa gtg agc cct tac agt gcc ccc tgg aaa ccc act 216
Cys Ala Val Gly Gln Val Ser Pro Tyr Ser Ala Pro Trp Lys Pro Thr
20 25 30
tgg cct gca tac cgc ctc cct gtc gtc ttg ccc cag tct acc ctc aat 264
10 Trp Pro Ala Tyr Arg Leu Pro Val Val Leu Pro Gln Ser Thr Leu Asn
35 40 45
tta gcc aag cca gac ttt gga gcc gaa gcc aaa tta gaa gta tct tct 312
Leu Ala Lys Pro Asp Phe Gly Ala Glu Ala Lys Leu Glu Val Ser Ser
50 55 60
15 tca tgt gga ccc cag tgt cat aag gga act cca ctg ccc act tac gaa 360
Ser Cys Gly Pro Gln Cys His Lys Gly Thr Pro Leu Pro Thr Tyr Glu
65 70 75
gag gcc aag caa tat ctg tct tat gaa acg ctc tat gcc aat ggc agc 408
Glu Ala Lys Gln Tyr Leu Ser Tyr Glu Thr Leu Tyr Ala Asn Gly Ser
20 80 85 90 95
cgc aca gag acg cag gtg ggc atc tac atc ctc agc agt agt gga gat 456
Arg Thr Glu Thr Gln Val Gly Ile Tyr Ile Leu Ser Ser Ser Gly Asp
100 105 110
ggg gcc caa cac cga gac tca ggg tct tca gga aag tct cga agg aag 504
25 Gly Ala Gln His Arg Asp Ser Gly Ser Ser Gly Lys Ser Arg Arg Lys
115 120 125
cgg cag att tat ggc tat gac agc agg ttc agc att ttt ggg aag gac 552

Arg Gln Ile Tyr Gly Tyr Asp Ser Arg Phe Ser Ile Phe Gly Lys Asp
130 135 140
ttc ctg ctc aac tac cct ttc tca aca tca gtg aag tta tcc acg ggc 600
Phe Leu Leu Asn Tyr Pro Phe Ser Thr Ser Val Lys Leu Ser Thr Gly
5 145 150 155
tgc acc ggc acc ctg gtg gca gag aag cat gtc ctc aca gct gcc cac 648
Cys Thr Gly Thr Leu Val Ala Glu Lys His Val Leu Thr Ala Ala His
160 165 170 175
tgc ata cac gat gga aaa acc tat gtg aaa gga acc cag aag ctt cga 696
10 Cys Ile His Asp Gly Lys Thr Tyr Val Lys Gly Thr Gln Lys Leu Arg
180 185 190
gtg ggc ttc cta aag ccc aag ttt aaa gat ggt ggt cga ggg gcc aac 744
Val Gly Phe Leu Lys Pro Lys Phe Lys Asp Gly Gly Arg Gly Ala Asn
195 200 205
15 gac tcc act tca gcc atg ccc gag cag atg aaa ttt cag tgg atc cgg 792
Asp Ser Thr Ser Ala Met Pro Glu Gln Met Lys Phe Gln Trp Ile Arg
210 215 220
gtg aaa cgc acc cat gtg ccc aag ggt tgg atc aag ggc aat gcc aat 840
Val Lys Arg Thr His Val Pro Lys Gly Trp Ile Lys Gly Asn Ala Asn
20 225 230 235
gac atc ggc atg gat tat gat tat gcc ctc ctg gaa ctc aaa aag ccc 888
Asp Ile Gly Met Asp Tyr Asp Tyr Ala Leu Leu Glu Leu Lys Lys Pro
240 245 250 255
cac aag aga aaa ttt atg aag att ggg gtg agc cct cct gct aag cag 936
25 His Lys Arg Lys Phe Met Lys Ile Gly Val Ser Pro Pro Ala Lys Gln
260 265 270
ctg cca ggg ggc aga att cac ttc tct ggt tat gac aat gac cga cca 984

Leu Pro Gly Gly Arg Ile His Phe Ser Gly Tyr Asp Asn Asp Arg Pro
275 280 285

ggc aat ttg gtg tat cgc ttc tgt gac gtc aaa gac gag acc tat gac 1032

Gly Asn Leu Val Tyr Arg Phe Cys Asp Val Lys Asp Glu Thr Tyr Asp
5 290 295 300

ttg ctc tac cag caa tgc gat gcc cag cca ggg gcc agc ggg tct ggg 1080

Leu Leu Tyr Gln Gln Cys Asp Ala Gln Pro Gly Ala Ser Gly Ser Gly
305 310 315

gtc tat gtg agg atg tgg aag aga cag cag cag aag tgg gag cga aaa 1128

10 Val Tyr Val Arg Met Trp Lys Arg Gln Gln Gln Lys Trp Glu Arg Lys
320 325 330 335

att att ggc att ttt tca ggg cac cag tgg gtg gac atg aat ggt tcc 1176

Ile Ile Gly Ile Phe Ser Gly His Gln Trp Val Asp Met Asn Gly Ser
340 345 350

15 cca cag gat ttc aac gtg gct gtc aga atc act cct ctc aaa tat gcc 1224

Pro Gln Asp Phe Asn Val Ala Val Arg Ile Thr Pro Leu Lys Tyr Ala
355 360 365

cag att tgc tat tgg att aaa gga aac tac ctg gat tgt agg gag ggg 1272

Gln Ile Cys Tyr Trp Ile Lys Gly Asn Tyr Leu Asp Cys Arg Glu Gly
20 370 375 380

tgacacag tgttccctcc tggcagcaat taagggtctt catgttctta ttttaggaga 1330

ggccaaattg tttttgtca ttggcgtgca cacgtgtgtg tgtgtgtgtg tgtgtaaggt 1390

gtcttataat cttttacctt tttcttacia ttgcaagatg actggcttta ctattgaaa 1450

actggtttgt gtatcatatc atatatcatt taagcagttt gaaggcatac tttgcatag 1510

25 aaataaaaaa aatactgatt tggggcaatg aggaatatt gacaattaag ttaattctca 1570

cgtttttgca aactttgatt tttatttcat ctgaacttgt ttcaaagatt tatattaaat 1630

atttggcata caagagatat gaattcttat atgtgtgcata gtgtgttttc ttctgagatt 1690

catcttggtg gtgggtttt ttgtttttt aattcagtc ctgacttta atgcttccat 1750
aaggcagtgt tcccatttag gaacttgac agcatttgtt aggagaata ttttgattt 1810
ggaggcattt gcatggtagt cttgaacag taaaatgatg tgttgactat actgatacac 1870
atattaaact ataccttata gtaaaccagt atcccaagct gcttttagtt caaaaaatag 1930
5 tttctttcc aaaggttgtt gctctacttt gtaggaagtc ttgcatatg gcctcccaa 1990
ctttaaagtc ataccagagt ggccaagagt gtttatcca acccttccat ttaacaggat 2050
ttactcaca ttctggaac tagctatttt tcagaagaca ataacaggg ctttaattaga 2110
acaggctgta ttctctcca gcaaacagtt gtggccacac taaaacaat catagcattt 2170
taccctgga ttatagcaca tctcatgttt tatcatttgg atggagtaat taaaatgaa 2230
10 ttaaattcca gagaacaatg gaagcattgc ctggcagatg tcacaacaga ataaccactt 2290
gtttggagcc tggcacagtc ctccagcctg atcaaaaatt attctgcata gtttcagtg 2350
tgctttctgg gagctatgta ctcttcaat ttggaaactt ttctctctca ttatagtga 2410
aaatacttgg aagtacttt aagaaaacca gtgtggcctt ttccctcta gctttaaag 2470
ggccgccttt gctggaatgc tctaggttat agataacaa ttaggtataa tagcaaaaat 2530
15 gaaaattgga agaattgcaa atggatcaga atcatgcctt ccaataaagg cctttacaca 2590
tgtttatca atatgattat caaatcacag catatacaga aaagacttgg acttattgta 2650
tgttttatt ttatggctct cggcctaagc acttcttctt aaatgtatcg gagaaaaaat 2710
caaatggact acaagcacgt gtttgctgtg cttgcacccc aggtaaacct gcattgtagc 2770
aatttgtaag gatattcaga tggagcactg tcacttagac attctctggg ggattttctg 2830
20 ctgtcttct ttgagctttt tggaaggata attctgataa ggcactcaag aaacgtacaa 2890
ccacagtgtt ttcttcaat catatgagaa atactatgca tagcaaggag atgcagagcc 2950
gccaggaaaa ttctgagttc cagcacaatt ttctttggaa tctaacagga atctagcctg 3010
aggaagaagg gaggtctcca ttctatgtc tggattttgg gggttttgtt tgttttct 3070
ttagcttggg gaaaaaaagt tctactgaaca ccaagaccag aatggatttt ttaaaaaaa 3130
25 tagatgttcc tttgtgaag caccttgatt ccttgatttt gattttttgc aaagttagac 3190
aatggcacia agtcaaaatg aaatcaatgt ttatgtcaca agtagatgta atttactaaa 3250
gaatgataca cccatatgct atatacagct taactcacag aactgtaaaa gaaaattata 3310

aaataattca acatgtccat ctttttagtg ataataaaag aaagcatggg attaaactat 3370
 catagaagta gacagaaaaa gaaaaaagga ctcattggcat tattaatata attagtgcct 3430
 tacatgtgtt agttatacat attagaagca tatttgccca gtaaggctag tagaaccaca 3490
 ttcccaaag tgtgctcctt aaacactcat gccttatgat ttctaccaa aagtaaaaag 3550
 5 ggtgttatta agtcagagga agatgcctct ccattttccc tcctttatc agagggtcac 3610
 atgcctgtct gcacattaaa agctctggga agacctgttg taaagggaca agttgaggtt 3670
 gtaaaatctg catttaaata aacatctttg atcac 3705

<210> 12
 10 <211> 383
 <212> PRT
 <213> Homo sapiens

 <400> 12
 15 Met Ala Gly Ile Pro Gly Leu Leu Phe Leu Leu Phe Phe Leu Leu
 1 5 10 15
 Cys Ala Val Gly Gln Val Ser Pro Tyr Ser Ala Pro Trp Lys Pro Thr
 20 25 30
 Trp Pro Ala Tyr Arg Leu Pro Val Val Leu Pro Gln Ser Thr Leu Asn
 20 35 40 45
 Leu Ala Lys Pro Asp Phe Gly Ala Glu Ala Lys Leu Glu Val Ser Ser
 50 55 60
 Ser Cys Gly Pro Gln Cys His Lys Gly Thr Pro Leu Pro Thr Tyr Glu
 65 70 75
 25 Glu Ala Lys Gln Tyr Leu Ser Tyr Glu Thr Leu Tyr Ala Asn Gly Ser
 80 85 90 95
 Arg Thr Glu Thr Gln Val Gly Ile Tyr Ile Leu Ser Ser Ser Gly Asp

100 105 110
 Gly Ala Gln His Arg Asp Ser Gly Ser Ser Gly Lys Ser Arg Arg Lys
 115 120 125
 Arg Gln Ile Tyr Gly Tyr Asp Ser Arg Phe Ser Ile Phe Gly Lys Asp
 5 130 135 140
 Phe Leu Leu Asn Tyr Pro Phe Ser Thr Ser Val Lys Leu Ser Thr Gly
 145 150 155
 Cys Thr Gly Thr Leu Val Ala Glu Lys His Val Leu Thr Ala Ala His
 160 165 170 175
 10 Cys Ile His Asp Gly Lys Thr Tyr Val Lys Gly Thr Gln Lys Leu Arg
 180 185 190
 Val Gly Phe Leu Lys Pro Lys Phe Lys Asp Gly Gly Arg Gly Ala Asn
 195 200 205
 Asp Ser Thr Ser Ala Met Pro Glu Gln Met Lys Phe Gln Trp Ile Arg
 15 210 215 220
 Val Lys Arg Thr His Val Pro Lys Gly Trp Ile Lys Gly Asn Ala Asn
 225 230 235
 Asp Ile Gly Met Asp Tyr Asp Tyr Ala Leu Leu Glu Leu Lys Lys Pro
 240 245 250 255
 20 His Lys Arg Lys Phe Met Lys Ile Gly Val Ser Pro Pro Ala Lys Gln
 260 265 270
 Leu Pro Gly Gly Arg Ile His Phe Ser Gly Tyr Asp Asn Asp Arg Pro
 275 280 285
 Gly Asn Leu Val Tyr Arg Phe Cys Asp Val Lys Asp Glu Thr Tyr Asp
 25 290 295 300
 Leu Leu Tyr Gln Gln Cys Asp Ala Gln Pro Gly Ala Ser Gly Ser Gly
 305 310 315

320 325 330 335

340 345 350

355 360 365

370 375 380

10